

QuestSpecialty

SHOOTER

Water-Based Fogger

FOR USE IN: Rooms, Apartments, Homes, Attics, Basements, Campers, Boats, Household Storage Areas, Garages, Pet Sleeping Areas and Cabins

One (6 ounce) canister of fogger (each can) treats a room up to 27 ft. x 27 ft. with an 8 ft. ceiling or 6,000 cubic feet of unobstructed space.

KILLS: Ants, Cockroaches, Fleas, Spiders, Mosquitoes, Houseflies & Others Listed

ACTIVE INGREDIENTS:

Tetramethrin	0.200%
(S)-cyano(3-phenoxyphenyl)methyl-(S)-4-chloro-alpha-(1-methylethyl)benzeneacetate	0.100%
* N-Octyl bicycloheptene dicarboximide	0.957%
OTHER INGREDIENTS	98.743%
	100.000%

* MGK® 264, Insecticide Synergist
EVERECIDE, MGK® - Registered trademarks of McLaughlin Gormley King Company

EPA REG. NO. 1021-1648-44446 EPA EST. NO. 44446-TX-1

For Spanish, refer to carton insert. Para español, refiérase a la adición del cartón.

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND FIRST AID ON BACK PANEL

NET WT. 6 OZ. (170 g)

Manufactured For
QuestSpecialty Corporation
P.O. Box 624 • Brenham, TX 77834
713.896.8188

600005440

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Breathing spray mist may be harmful. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.



PHYSICAL OR CHEMICAL HAZARDS Highly Flammable Ingredient Ingrediente Altamente Inflamable

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. This product contains a highly flammable ingredient. It may cause a fire or explosion if not used properly. Follow the Directions for Use on this label very carefully.

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.

This product will kill ants, cockroaches, confused flour beetles, crickets, fleas, houseflies, mosquitoes, rice weevils, saw-toothed grain beetles, small flying moths, saw bugs, spiders (including black widow), and brown dog ticks.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE

Does not control bedbugs.

GENERAL PRECAUTIONS AND RESTRICTIONS: Do not apply this product in a manner that will contact adults, children or pets either directly or through drift. Remove pets, birds, and cover fish aquariums before spraying. Do not allow adults, children, or pets to enter the treated area, until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated. For use only when building has been vacated by human beings and pets. Wait two (2) hours after application, then open windows, vents and doors for two (2) hours. If an odor is still detectable, continue ventilating until no remaining odor of pesticide remains.

TO USE THIS PRODUCT CORRECTLY: DO NOT use more than one fogger per room. DO NOT use in small, enclosed spaces such as closets, cabinets or under counters or tables.

Use of a fogger in an enclosed space may cause the product to explode, resulting in injury to people or damage to property.



Do not use in a room 5 ft. X 5 ft. or smaller; instead, allow fog to enter from other rooms. Turn off ALL ignition sources such as pilot lights (shut off gas valves), other open flames or running electrical appliances that cycle off and on (i.e., refrigerators, thermostats, etc.). Call your gas utility or management company if you need assistance with your pilot lights.



Use only one canister of fogger to treat a room up to 27 ft. x 27 ft. with an 8 ft. ceiling or 6,000 cubic feet of unobstructed space.

BEFORE YOU FOG: In the home do not apply when food is present. Exposed food should be covered or removed. All food preparation surfaces and utensils should be covered during treatment or thoroughly washed before use.

Close outside doors and windows. Shut off fans, air conditioners, and ventilation equipment. Put out all flames, pilot lights and other ignition sources. Call your gas utility or management company if you need assistance.

Remove pets and birds, cover and turn off fish aquariums before activating fogger. Place a chair, small table or stand in the center of the room to be treated and cover with newspaper. Cover wood floors and wood furniture in the immediate area around fogger with newspapers or a drop cloth. Open cabinets, cupboards, drawers, closets and doors in areas to be treated. Fill out and place hang-tag on the door to the treated area to alert family and others with access to the treated area not to enter for 2 hours.

SHAKE WELL BEFORE USE. KEEP CONTAINER UPRIGHT

TO START FOGGING: Point fogger away from face and eyes when releasing (container sprays straight up). To lock valve in open position for automatic discharge, press valve button down firmly, until it snaps into place. Then place the fogger on the stand or table in the center of the room.

Vacate the treated house, individual apartment unit or other structure immediately. Do not remain in the area during treatment. Keep treated area closed for two (2) hours.

Do not Re-enter for 2 hours

AIRING OUT: Wait two (2) hours after application, then open windows, vents and doors for two (2) hours. If an odor is still detectable, additional ventilation is required.

To Kill Brown Dog Ticks and Fleas: Destroy pet's old bedding and replace with clean, fresh bedding after treatment is completed. Use product as above. Pets should be treated with a registered flea and tick control product prior to reentering treated premises.



STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **Pesticide Storage:** Store in a cool, dry area. Always store pesticides in the original container. Store away from food or pet food. **Pesticide Disposal and Container Handling:** Do Not Puncture or Incinerate! Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions.



ITEM 4400