

# Country Carden Hospital Disinfectant Decodorant

Virucidal\* • Tuberculocidal \*\*\*\* Fungicidal\*\*\* • Bactericidal\*\* Pseudomonacidal Against P. aeruginosa Staphylocidal Against S. aureus Sanitizes Fabric and Leather Controls Mold/Mildew Fungi for 7 Days On Cotton, Fabric & Leather

(Specific species/agents listed on back panel.)

PLEASANTLY SCENTED • KILLS MRSA\*\*

\*KILLS AVIAN INFLUENZA A VIRUS H5N1

\*KILLS HIV-1 (AIDS VIRUS) AND HERPES SIMPLEX TYPES 1 & 2 VIRUSES ON PRECLEANED **ENVIRONMENTAL SURFACES/OBJECTS** PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS.

FOR USE ON PRECLEANED, HARD, NONPOROUS, NONFOOD-CONTACT SURFACES IN/ON INDUSTRIAL, INSTITUTIONAL, COMMERCIAL, MEDICAL, AND RESIDENTIAL FACILITIES, EQUIPMENT, AND VEHICLES,

Authorized by USDA for use in Federally inspected meat and poultry plants.

ACTIVE INGREDIENTS:	ACT	VE	ING	RED	IEN	TS:
---------------------	-----	----	-----	-----	-----	-----

2-Phenylphenol			0.209
Ethanol			53.009
INERT INGREDIENTS			46.809
TOTAL	FIL Famous	ALCOHOL:	100 009

## KEEP OUT OF REACH OF CHILDREN WARNING:

See additional precautionary and first aid statements on back panel.

NET WT. 16.5 OZ. (468 g)

QuestSpecialty Corporation P.O. Box 624 • Brenham, TX 77834 • 713.896.8188

#### FEATURES, CLAIMS, AND USES

This product is an effective Virucide\*, Bactericide\*\*, Tuberculocide \*\*\*\* and Fungicide\*\*\* on hard nonporous surfaces when used as a disinfectant. This product has demonstrated effectiveness on precleaned hard nonporous surfaces against the following microorganisms:

VIRUCIDAL\* - Kills Human Immunodeficiency virus Type 1 (HIV-1) Virus, Herpes Simplex Virus Type 1 and Herpes Simplex Virus Type 2, Influenza A2/H3N2 Virus, Adenovirus Type 5, Vaccinia Virus, Infectious Canine Adenovirus 1, Canine Distemper Virus, and Avian Influenza A Virus (H5N1), \*SARS-CoV-2 virus the cause of \*COVID-19. Follow instructions for disinfection

BACTERICIDAL\*\* - Kills: Escherichia coli; Pseudomonas aeruginosa; Salmonella enterica; Salmonella schottmuelleri (paratyphoid B); Shigella dysenteriae; Streptococcus pyogenes; Staphylococcus aureus; and Methicillin-resistant Staphylococcus aureus (MRSA) Follow instructions for disinfection.

TUBERCULOCIDAL\*\*\*\* - Kills Mycobacterium tuberculosis bovis

BCG (Tubercle bacilli). Follow instructions for disinfection.

FUNGICIDAL\*\*\* - Kills Trichophyton interdigitale ("athlete's foot" fungi; formerly known as T. mentagrophytes) in bathrooms, shower areas, and locker rooms. Follow instructions for disinfection.

CONTROLS MOLD/MILDEW - Inhibits the growth of Aspergillus niger and Penicillium glaucum fungi on fabric for up to 7 days. Repeat treatment as necessary

SANITIZES - Controls Staphylococcus aureus and Klebsiella pneumoniae on porous cotton and polyester and leather surfaces, such as on shoes, athletic mats, athletic shoes and other athletic equipment. For use as a spot treatment.

DEODORIZES - Pleasant fragrance quickly suppresses existing unpleasant odors from many sources.

FOR USE IN: Industrial, Institutional, Commercial, Medical, and Residential facilities, equipment, and vehicles, including hospitals; ambulances; nursing homes; medical and dental offices and clinics; veterinary offices and animal areas; schools; laboratories; hotels and motels; restaurants and cafeterias; food storage, processing, packaging, handling, and serving establishments and equipment; offices; stores; factories and manufacturing plants; and apartment buildings and homes.

FOR USE ON: Most glass, glazed porcelain, glazed ceramic, enameled, painted, plastic (such as ABS, acrylic, latex rubber, phenolic resin, polyamide, polycarbonate, polyester, polyethylene, polyolefin, polypropylene, polystyrene, polytetrafluorethylene, polyurethane, and vinyl), rubber, metal, sealed leather, finished wood, sealed grout and sealed concrete, and similar nonporous surfaces.

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Harmful if absorbed through skin. Do not get in eyes or on skin or clothing. Wear protective eyewear such as goggles, face shield or safety glasses. Wear long-sleeved shirt and long pants, shoes and socks, and chemical resistant gloves (such as or made out of any water proof material). Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

#### FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a Poison Control Center or doctor for treatment advice

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. For general information on product use, etc. call the National Pesticides Information Center at 1-822-858-7378, You may also contact the poison control center at 1-800-222-1222 for emergency medical treatment information.

PHYSICAL/CHEMICAL HAZARDS: Flammable. Contents under pressure. Keep away from heat, sparks, pilot lights, and flames. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54°C) may cause bursting.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner incon-

infectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

GENERAL: Shake container well before each use and hold it upright while spraying. For use on precleaned, hard nonporous surfaces; not for use on humans or animals. Not for use where food or related products are prepared, processed or stored. Remove food or related products from the area being treated. Avoid use where product would contact counters or tabletops where food may be stored, prepared or processed. Do not use on waxed/polished surfaces, water-based "latex" paint, shellac, cellulose (such as rayon) or other alcohol-sensitive materials. Clean surfaces to be treated with a suitable hard surface cleaner removing visible soils. Allow to air dry prior to treatment.

To disinfect hard, non-porous surfaces: Preclean surfaces as directed above. Hold container upright about 6-8" from surface. Spray to thoroughly wet hard, nonporous surfaces. Treated surfaces must remain visibly wet for 10 minutes. Allow to air dry. Rinse all surfaces that come in contact with food such as counters, appliances, tables with potable water. Do not use on utensils, glassware, and dishes. For \*SARS-CoV-2, treated surfaces must remain visibly wet for 3 minutes.

Before use in federally inspected meat and poultry plants, all food products and packaging materials must be removed or carefully protected before disinfection

To disinfect toilets and urinals: Remove visible soil prior to disinfection. Empty toilet bowl or urinal. Hold container upright about 6-8" from surface. Spray to thoroughly wet hard, nonporous surfaces above the water line. Treated surfaces must remain visibly wet for 10 minutes. Brush over exposed surfaces. Allow to stand for 10 minutes, then flush.

To disinfect Veterinary offices, animal areas, and animal laboratories: Remove all animals and feed from premises, vehicles, cages, kennels, etc. Remove all litter, droppings, and manure from floors, walls, and surfaces of facilities occupied or traversed by animals. Empty all feeding and water appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Hold container upright about 6-8" from surface. Spray to thoroughly wet hard, nonporous surfaces. Treated surfaces must remain wet for 10 minutes. Allow to air dry, Immerse all halters, collars, leashes and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, vehicles, and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub feeding and watering equipment with soap or detergent, and rinse with potable water before reuse.

To Clean/Deodorize: Hold container upright about 6-8" from surface. Spray surface and allow to air dry.

To Control Mold/Mildew: For use as a spot treatment. Inhibits the growth of Aspergillus niger and Penicillium glaucum fungi on porous surfaces such as cotton and polyester fabric and leather surfaces. Hold container upright about 6-8" from surface. Spray to thoroughly wet surfaces. Treated surfaces must remain visibly wet for 5 minutes. Allow surfaces to air dry. For mold/mildew control, repeat application every 7 days or more frequently if new growth appears.

To Sanitize Cotton or Polyester Surfaces: For use as a spot treatment. Controls Staphylococcus aureus and Klebsiella pneumoniae on porous surfaces such as cotton and polyester fabric and leather surfaces. Hold container upright about 6-8" from surface. Spray to thoroughly wet surfaces. Treated surfaces must remain visibly wet for 5 minutes. Allow surfaces to air dry.

\*TO KILL HIV-1 (Human Immunodeficiency virus Type 1; AIDS virus) ON PRE-CLEANED HARD NON-POROUS SURFACES/OBJECTS in health care or other settings with inanimate environmental surfaces/objects associated with the potential for transmission of HIV-1 and likely to be soiled with blood/body fluids, follow these additional SPECIAL INSTRUCTIONS FOR CLEANING AND DE-CONTAMINATING AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: PERSONAL PROTECTION - Barrier protection items such as disposable latex gloves, gowns, masks, and eye coverings must be worn when handling items soiled with blood/body fluids. CLEANING PROCE-DURE - Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of this product as a disinfectant. CONTACT TIME - 10 minutes. DISPOSAL OF INFECTIOUS MATERIALS - Blood/body fluids and other materials removed from or used to clean surfaces/objects must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. Treated surfaces must remain visibly wet for 10 minutes.

#### STORAGE AND DISPOSAL

GENERAL: Do not contaminate other materials (including foods/drinks/feeds/water) during transport, use, storage, and disposal. If damaged or leaking, soak up and wrap any waste, then dispose of as below.

STORAGE: Keep in a cool, dry, locked area inaccessible to children and away from sources of heat and ignition (including flames, sparks, hot surfaces, and sunlight).

PESTICIDE DISPOSAL: If empty: Place in trash or offer for recycling if available. If partially filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. CONTAINER HANDLING: Replace protective cap on container. Wrap container in several layers of paper and place in trash or offer for recycling if available. Do not puncture or incinerate.

HMIS Ratings: Health - 1. Flammability - 3. Reactivity - 1. Personal Protection - A.

ITEM 311500001-20AR

Made in USA

7 8 8

\_ ღ

**=** ∞

0

